Emetrol Drug Facts

Drug Facts	
Active ingredient (per 5mL)	Purpose
Dextrose (glucose) 1.87g	Upset stomach reliever
Levulose (fructose) 1.87g	Upset stomach reliever
Phosphoric acid 21.5mg	Upset stomach reliever

Uses

• for relief of upset stomach associated with nausea.

Warnings

■ this product contains fructose and should not be taken by persons with hereditary fructose intolerance (HFI)

Do not use if you have

■ allergic reactions to any of the ingredients in this product

Ask a doctor before use if you have ■ diabetes

Stop use and ask a doctor if

■ symptoms persist, return or get worse

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Directions

- for maximum effectiveness never dilute or drink fluids of any kind immediately before or after taking this product
- repeat dose every 15 minutes or until distress subsides
- do not take more than 5 doses in 1 hour without consulting a doctor
- measure only with dosing cup provided. Dosing cup to be used with this product only. Do not use with other products.
- mL = milliliters

age	dose
Adults and children 12 years of age and over	15 mL or 30 mL
Children 2 to under 12 years of age	5 mL or 10 mL

Other information

- store between 15°-30°C (59°-86°F) away from heat and direct light; keep from freezing.
- do not use if printed seal under cap is broken or missing

Inactive ingredients FD&C red no. 40, flavors, glycerin, methylparaben, purified water

Questions or comments? 1-844-241-5454