

Emetrol Drug Facts

| Drug Facts | |
|---|------------------------|
| Active ingredient (per 5mL) | Purpose |
| Dextrose (glucose) 1.87g | Upset stomach reliever |
| Levulose (fructose) 1.87g | Upset stomach reliever |
| Phosphoric acid 21.5mg | Upset stomach reliever |
| Uses | |
| <ul style="list-style-type: none"> ■ for relief of upset stomach associated with nausea. | |
| Warnings | |
| <ul style="list-style-type: none"> ■ this product contains fructose and should not be taken by persons with hereditary fructose intolerance (HFI) | |
| Do not use if you have | |
| <ul style="list-style-type: none"> ■ allergic reactions to any of the ingredients in this product | |
| Ask a doctor before use if you have ■ diabetes | |
| Stop use and ask a doctor if | |
| <ul style="list-style-type: none"> ■ symptoms persist, return or get worse | |
| If pregnant or breast feeding , ask a health professional before use. | |
| Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. | |
| Directions | |
| <ul style="list-style-type: none"> ■ for maximum effectiveness never dilute or drink fluids of any kind immediately before or after taking this product ■ Adults and children 12 years of age and over : 15mL or 30mL ■ Children 2 to under 12 years of age: 5mL or 10mL ■ repeat dose every 15 minutes or until distress subsides ■ do not take more than 5 doses in 1 hour without consulting a doctor ■ measure only with dosing cup provided. Dosing cup to be used with this product only. <p style="margin-left: 20px;">Do not use with other products.</p> <p>mL = milliliters</p> | |
| Other information | |
| <ul style="list-style-type: none"> ■ store between 15°-30°C (59°-86°F) away from heat and direct light; keep from freezing. ■ do not use if printed seal under cap is broken or missing | |
| Inactive ingredients FD&C red no. 40, flavors, glycerin, methylparaben, purified water | |
| Questions or comments? 1-844-241-5454 | |