

Emetrol Chewables

Drug Facts	
Active ingredient (in each chewable tablet)	Purpose
Sodium citrate dihydrate USP 230 mg	Upset stomach reliever/antacid
Uses For the relief of nausea associated with upset/sour stomach, including that due to overindulgence in food and drink.	
Warnings	
Do not use if you ■ have Hereditary Fructose Intolerance (HFI). This product contains fructose.	
Ask a doctor before use if you ■ have diabetes because this product contains sugar ■ are on a sodium-restricted diet	
Ask a doctor or pharmacist before use if you are taking any other medications. This product may interact with certain prescription drugs.	
When using this product, do not take more than 24 tablets in a 24-hour period.	
Stop use and ask a doctor if nausea lasts more than two weeks or recurs frequently.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
■ Adults: 2-4 tablets. ■ Children: Consult a doctor for appropriate dosage.	
■ Chew tablets completely. Do not swallow whole ■ Dosage may be repeated after 15 minutes, not to exceed 24 tablets in a 24-hour period unless advised by a doctor. ■ Read all package directions and warnings before use and use only as directed. ■ These tablets are intended for use by normally healthy persons only.	
■ Persons under 18 years of age should use only as directed by a doctor.	
Other information	
■ Each tablet contains: sodium 60 mg ■ Store at room temperature ■ Protect product from excessive heat, cold and moisture. ■ Note: This product is not intended as a substitute for a balanced nutritional diet or as an electrolyte replenishment. ■ Do not use if printed seal under cap is broken or missing.	
Inactive ingredients dextrose, flavor, fructose, hydroxypropyl cellulose, magnesium stearate, maltodextrin, silica, sodium carboxymethylcellulose, sucralose.	
Questions or comments? 1-844-241-5454 or Emetrol.com	